

Preprint submitted to

Acta Phys. Hung. A 22/1 (2005) 000–000

HEAVY ION
PHYSICS

Dissipation and fragmentation of low- Q^2 scattered partons in Au-Au collisions at RHIC

R. L. Ray and M. Daugherty, STAR Collaboration

Department of Physics, The University of Texas at Austin,
Austin, Texas 78712, USA

Received 1 January 2005

Abstract. Two-particle correlations and event-wise fluctuations in transverse momentum p_t are reported for Au-Au collisions at $\sqrt{s_{NN}} = 62$ and 200 GeV on pseudorapidity η and azimuth ϕ . Distributions of all pairs of particles (no leading trigger particle) reveal jet-like correlations, or peaks at pair-wise opening angles of order 1 radian or less. The width of this *same-side* correlation peak increases dramatically on pseudorapidity and decreases on azimuth for increasing collision centrality. Evolution of the same-side peak with centrality suggests dissipation of low- Q^2 partons via strong coupling to an expanding bulk medium. p_t correlations, which provide access to temperature and/or velocity distributions in the colliding system, are also presented.

Keywords: correlations, fluctuations, Au-Au collisions, minijets

PACS: 24.60.Ky, 25.75.Gz

1. Introduction

Theoretical descriptions of relativistic heavy-ion collisions predict abundant low- Q^2 gluon production in the early stages of the collision with rapid parton thermalization driving the formation of a colored medium [1, 2, 3]. Remnants of these semi-hard processes may survive to the final, decoupling stage owing to the finite size and lifetime of the collision volume. If so, there should be contributions to nonstatistical event-wise fluctuations of mean- p_t $\langle p_t \rangle$ and two-particle correlations. Large non-statistical $\langle p_t \rangle$ fluctuations have been reported for Au-Au collisions at 130 GeV [4] and 200 GeV [5] which are much larger than was observed at the SPS [6, 7]. Study of the related two-particle correlations facilitates interpretation of event-wise fluctuations in terms of underlying dynamics. In this paper we report preliminary two-particle correlations for Au-Au collisions at 62 GeV on (η, ϕ) and $\langle p_t \rangle$ fluctua-

tions and p_t correlations for Au-Au at 200 GeV [8] from the STAR Collaboration. Low- Q^2 partons are emphasized in order to probe nonperturbative QCD medium effects. These data complement the two-particle correlations for Au-Au collisions at 130 GeV reported in [9, 10].

Data for the present analysis were obtained with the STAR detector [11] where charged particles were accepted in $|\eta| \leq 1.3$, 2π azimuth, and $p_t \geq 0.15$ GeV/ c . Corrections were made for two-track inefficiencies and charge-sign dependent cuts were applied to minimize quantum and Coulomb correlation contributions. These cuts do not significantly affect the correlation structures shown here. Cuts based on dE/dx in the STAR Time Projection Chamber tracking detector were applied to reduce photon conversion $e^\pm$ contamination. See Ref. [10] for discussion of the various cuts.

2. Autocorrelations

In conventional jet analysis event-wise concentrations of transverse momentum or energy localized on angle variables (η, ϕ) are identified. In heavy ion collisions, where such identification is impractical, jet studies are based on a high- p_t ‘leading particle’ which may estimate a parton momentum direction and some fraction of its magnitude. However, for low- Q^2 partons identification of a ‘leading particle’ is not possible. Furthermore the number of correlated hadrons from each low- Q^2 parton is expected to be relatively small while the number of such partons per event is large (of order tens). Observation of these relatively soft correlated hadrons is accessed via *autocorrelations* which are well-known in signal processing disciplines to enable weak but repetitive signals, which cannot be detected individually, to be accurately measured in aggregate. An autocorrelation is a projection *by averaging* from subspace (x_1, x_2) onto difference variable $x_\Delta \equiv x_1 - x_2$. The autocorrelations reported in [9, 10] and here do not require a leading- or trigger-particle, but instead use all particle pairs within the acceptance.

The two-particle correlation density per final state particle is defined by

$$\Delta\rho/\sqrt{\rho_{mix}} \equiv [\rho_{sib}(\vec{p}_1, \vec{p}_2) - \rho_{mix}(\vec{p}_1, \vec{p}_2)]/\sqrt{\rho_{mix}}, \quad (1)$$

where $\rho_{sib}(\vec{p}_1, \vec{p}_2)$ is the object distribution comprised of particle pairs from single events (*sibling* pairs) and the reference distribution, $\rho_{mix}(\vec{p}_1, \vec{p}_2)$, consists of pairs formed by sampling each particle of the pair from two different but similar events (*mixed-event* pairs). Studies of measured two-particle correlation projections onto subspaces $(p_{t1}$ vs $p_{t2})$, $(\eta_1$ vs $\eta_2)$ and $(\phi_1$ vs $\phi_2)$ indicate that the principle dependences for Au-Au collision data are on $(p_{t1} + p_{t2})$, $(p_{t1} - p_{t2})$, and the differences $\eta_\Delta \equiv \eta_1 - \eta_2$ and $\phi_\Delta \equiv \phi_1 - \phi_2$. For the latter two the correlation distributions are simultaneously projected onto *difference variables* ϕ_Δ and η_Δ ; the projection is then referred to as a *joint autocorrelation*.

3. Charge-Independent Joint Autocorrelations on $(\eta_\Delta, \phi_\Delta)$

Plotted in Fig. 1 are perspective views of preliminary charge-independent joint autocorrelations $\Delta\rho/\sqrt{\rho_{mix}}$ on difference variables $(\eta_\Delta, \phi_\Delta)$ for three centrality bins (approximately 80-90%, 50-60% and 10-20%) for Au-Au collisions at 62 GeV. The distributions are dominated by 1) a $\cos(2\phi_\Delta)$ component attributed to elliptic flow; 2) a $\cos(\phi_\Delta)$ component associated with transverse momentum conservation, and 3) a 2D same-side ($|\phi_\Delta| < \pi/2$) peak, which is the principal object of interest here, and assumed to be associated with low- Q^2 scattered parton fragmentation. In addition, for the most-peripheral data, a ϕ_Δ independent gaussian distribution on η_Δ is observed similar to that seen in correlation studies of p-p collision data [12] and assumed to be due to charge-ordering associated with longitudinal string fragmentation. This feature vanishes quickly with centrality. The same-side peak




Fig. 1. $\Delta\rho/\sqrt{\rho_{ref}}$ for Au-Au collisions at $\sqrt{s_{NN}} = 62$ GeV on $(\eta_\Delta, \phi_\Delta)$ for centralities ~ 80 -90%, 50-60% and 10-20% fraction of Au-Au total cross section.

in Fig. 1 varies strongly with centrality, transitioning from nearly symmetric on $(\eta_\Delta, \phi_\Delta)$ for peripheral collisions to dramatically broadened along η_Δ and narrowed on ϕ_Δ for the more central collisions. Resonance (*e.g.* ρ^0, ω) decays contribute about 3% of the peaks at (0,0) [10]. Electron conversion pairs which remain after the dE/dx cuts add to the bin at (0,0) producing the narrow spike there.

Mean- p_t fluctuation measure $\Delta\sigma_{p_t:n}^2(\delta\eta, \delta\phi)$ [4] obtained for particles within a two-dimensional bin of size $(\delta\eta, \delta\phi)$, is equivalent to an integral of p_t autocorrelation $\Delta\rho(p_t : n; \eta_\Delta, \phi_\Delta)/\sqrt{\rho_{mix}}$ over difference variables from (0,0) to $(\delta\eta, \delta\phi)$ [8]. Inversion of this integral equation yields p_t autocorrelations. The latter provide access to temperature and/or velocity distributions in the colliding system independent of that afforded by number of pair correlations [8]. Fig. 2 shows $\Delta\sigma_{p_t:n}^2(\delta\eta, \delta\phi)$ (left panel), the resulting p_t autocorrelation (middle panel), and p_t autocorrelation with $\cos(2\phi_\Delta)$ and $\cos(\phi_\Delta)$ subtracted [8] (right panel). The same-side peak structure broadens dramatically with centrality (not shown) on η_Δ as in Fig. 1 but remains narrower than the corresponding number of pair correlations.




Fig. 2. $(\delta\eta, \delta\phi)$ scale-dependent $\langle p_t \rangle$ fluctuation, p_t autocorrelation, and p_t auto-correlation with $\cos(2\phi_\Delta)$ and $\cos(\phi_\Delta)$ terms subtracted for mid-central (20-30%) Au-Au collisions at 200 GeV. ρ_{ref} is the same as ρ_{mix} in the text.

4. Conclusions

In STAR we are studying how the same-side autocorrelation structures evolve with centrality in Au-Au collisions at $\sqrt{s_{NN}} = 62, 130$ and 200 GeV. Same-side charge-average pair number and p_t autocorrelations strongly elongate on η and narrow on ϕ . The STAR experiment results reported here and elsewhere [8, 9, 10] are consistent with a picture in which (1) semi-hard parton scattering transfers momentum (p_t) to the bulk medium, inducing temperature/velocity fluctuations in the soft p_t range whose correlation amplitudes scale with the number of binary collisions per participant, and (2) particles correlated with semi-hard scattered partons in the early stages of the collision interact with the longitudinally expanding medium resulting in larger average $|\eta_\Delta|$. The reduced width on ϕ is not understood.

References

1. K. Kajantie, P. V. Landshoff and J. Lindfors, *Phys. Rev. Lett.* **59** (1987) 2527.
2. A. H. Mueller, *Nucl. Phys. B* **572** (2000) 227.
3. G. C. Nayak, A. Dumitru, L. D. McLerran and W. Greiner, *Nucl. Phys. A* **687** (2001) 457.
4. J. Adams *et al.* (STAR Collaboration), *Phys. Rev. C* **71** (2005) 064906.
5. S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **93** (2004) 092301.
6. D. Adamová *et al.* (CERES Collaboration), *Nucl. Phys. A* **727** (2003) 97.
7. T. Anticic *et al.* (NA49 Collaboration), *Phys. Rev. C* **70** (2004) 034902.
8. J. Adams *et al.* (STAR Collaboration), eprint nucl-ex/0509030, submitted to *Phys. Lett. B*; T. Trainor, R. Porter and D. Prindle, *J. Phys. G: Nucl. Part. Phys.* **31** (2005) 809.
9. J. Adams *et al.* (STAR Collaboration), eprint nucl-ex/0408012.
10. J. Adams *et al.* (STAR Collaboration), eprint nucl-ex/0411003.
11. K. H. Ackermann *et al.*, *Nucl. Instrum. Meth. A* **499** (2003) 624; see other STAR papers in volume **A499**.
12. R. J. Porter and T. A. Trainor (STAR Collaboration), eprint hep-ph/0406330.